

## REVIEW ARTICLE

# Skin-to-skin contact the first hour after birth, underlying implications and clinical practice

Ann-Marie Widström<sup>1</sup> , Kajsa Brimdyr<sup>2</sup> , Kristin Svensson<sup>1,3</sup> , Karin Cadwell<sup>2</sup> , Eva Nissen (eva.nissen@ki.se)<sup>1</sup> 

1. Karolinska Institutet, Stockholm, Sweden

2. Healthy Children Project Inc., Sandwich, MA, USA

3. Karolinska University Hospital, Karolinska Institutet, Stockholm, Sweden

## Keywords

Breastfeeding, Clinical practice, Maternal behaviour, Newborn infant behaviour, Skin-to-skin contact

## Correspondence

Eva Nissen, Karolinska Institutet, Widerströmska Huset, Tomtebodavägen 18 a, 171 77 Stockholm, Sweden.

Tel: +46737121553 |

Fax: +46737121553 |

Email: eva.nissen@ki.se

## Received

20 April 2018; revised 28 January 2019; accepted 11 February 2019.

DOI:10.1111/apa.14754

## INTRODUCTION

The evidence supporting the practice of skin-to-skin contact after birth is robust, indicating multiple benefits for both mother and baby. The 2016 Cochrane Review supports using immediate or early skin-to-skin contact to promote breastfeeding (1). Advantages for the mother include earlier expulsion of the placenta (2,3), reduced bleeding (3), increased breastfeeding self-efficacy (4) and lowered maternal stress levels (5). It has been suggested that the rise in the mother's oxytocin during the first hour after birth is related to the establishment of mother–infant bonding (6, 7). Advantages for the baby include a decrease of the negative consequences of the 'stress of being born' (8), more optimal thermoregulation (9), continuing even in the first days (8) and less crying (10). Skin-to-skin contact has been shown to increase breastfeeding initiation and exclusive breastfeeding while reducing formula supplementation in hospital, leading to an earlier successful first breastfeed (2,11,12), as well as more optimal suckling (3,13). The evidence supporting skin-to-skin the first hour is so compelling that the 2018 revision of the World Health Organization (WHO)/United Nations International Children's Emergency Fund (UNICEF) *Ten Steps to Successful Breastfeeding* that form the basis of the Baby-Friendly

## ABSTRACT

**Aim:** This paper integrates clinical expertise to earlier research about the behaviours of the healthy, alert, full-term infant placed skin-to-skin with the mother during the first hour after birth following a noninstrumental vaginal birth.

**Method:** This state-of-the-art article forms a link within the knowledge-to-action cycle, integrating clinical observations and practice with evidence-based findings to guide clinicians in their work to implement safe uninterrupted skin-to-skin contact the first hours after birth.

**Results:** Strong scientific research exists about the importance of skin-to-skin in the first hour after birth. This unique time for both mother and infant, individually and in relation to each other, provides vital advantages to short- and long-term health, regulation and bonding. However, worldwide, clinical practice lags. A deeper understanding of the implications for clinical practice, through review of the scientific research, has been integrated with enhanced understanding of the infant's instinctive behaviour and maternal responses while in skin-to-skin contact.

**Conclusion:** The first hour after birth is a sensitive period for both the infant and the mother. Through an enhanced understanding of the newborn infant's instinctive behaviour, practical, evidence-informed suggestions strive to overcome barriers and facilitate enablers of knowledge translation. This time must be protected by evidence-based routines of staff.

Hospital Initiative. Step 4 states 'Facilitate immediate and uninterrupted skin-to-skin contact and support mothers to initiate breastfeeding as soon as possible after birth' (14) with the understanding that the newborn infant will self-attach.

During this first hour after childbirth, both mother and newborn infant experience a special and unique time, a sensitive period (15,16), which has been biologically predetermined, especially after vaginal birth. This is aided by

## Key notes

- A lag exists between research knowledge and clinical practice surrounding skin-to-skin in the first hour after birth.
- A mutual early sensitive period includes preprogrammed behaviours for bonding and other survival mechanisms, for example through the eye contact, suggesting long-term implications.
- The framework of the newborn infant's 9 stages creates an opportunity to understand the biological and physiological situation for the dyad, and clinical implications of practice during this sensitive time.

the physiological state of each: the mother's high oxytocin levels and newborn infant's extremely high catecholamine levels.

Being in skin-to-skin contact with the mother after birth elicits the newborn infant's internal process to go through what could be called 9 *instinctive stages*: birth cry, relaxation, awakening, activity, rest, crawling, familiarization, suckling and sleeping (17) (Table 1).

This process is suggested to contribute to an early coordination of infant's five senses: sight, hearing, touch, taste and smell, as well as movement (17,18).

Whereas prolactin is the most important milk-producing hormone, oxytocin plays a key role in maternal behaviour and bonding immediately after birth. In animal studies, for example, if an ewe is separated from her lamb soon after birth, she will reject the lamb when they later are reunited. Interestingly, it has been shown that simulating a birth through the birth channel at reunion of mother and lamb enhances oxytocin release in the mother ewe, resulting in her acceptance of her lamb. This illustrates that natural oxytocin facilitates the bonding between the ewe and the lamb during the first hour after birth (19).

In the human mother, a surge of oxytocin is released in the mother's blood vessels during the first hour after birth to contract the uterus, facilitate placental discharge and to decrease blood loss (6). Oxytocin released to the blood stream in this situation is likely to be paralleled by an intense firing of parvocellular oxytocin neurons in the brains, as Theodosius has shown in animal experiments (20), causing an increased maternal sensitivity for the young. In humans, this has been illustrated through the mother's desire to keep her infant close to her throughout the hospital stay if the newborn infant suckled or even just touched her nipple during the first hour while skin-to-skin (21).

The mother is attracted to the infant's smell, facilitating a chemical communication between the two (22,23). This

highlights the importance of a new mother's access to her newborn infant's bare head to smell her baby. This is an example of the early symbiotic biological relationship between the dyad (24,25).

The newborn infant has high levels of catecholamine immediately after a normal, vaginal birth (10). These are highest closest to the time of birth, especially for the first thirty minutes. Catecholamines strengthen memory and learning.

In newborn infants less than 24 hours old, the odour of the mother's colostrum increases the amount of oxygenated haemoglobin over the olfactory cortex, as measured by near-infrared spectroscopy. In addition, the closer to birth, the more oxygenated haemoglobin was observed over the olfactory cortex (26). This increased sensitivity for the odour of breastmilk, especially soon after the birth, indicates a physiologically based early sensitive period in the newborn infant. This reaction matches the mother's biologically based enhancements of the breast odour through the increase of the surface of the areola and Montgomery gland secretions during the corresponding time (27). Thus, the increased early sensitivity for the odour of breastmilk, in the presence of high levels of catecholamine which strengthens this memory, is indication of a physiologically based early sensitive period in the newborn infant.

### Aim

The aim of this paper is to integrate clinical expertise to earlier research about the behaviours of the healthy, alert, full-term infant placed skin-to-skin with the mother during the first hour after birth following a noninstrumental vaginal birth.

### METHOD

This is a state-of-the-art article integrating clinical observations and practice with evidence-based findings to guide clinicians in their work to implement safe uninterrupted skin-to-skin contact the first hours after birth. This paper, presented from the nurse clinician perspective, is informed by the scientific literature around newborn infant behaviour and mother-to-infant interaction, in order to translate existing knowledge and facilitate clinician behaviour change. In this way, it forms a link within the knowledge-to-action cycle and offers the underlying implications of this important behaviour change to enhance the translation between the evidence and implementation through a deeper understanding of the newborn infant's instinctive behaviour while skin-to-skin immediately after birth.

This paper is organised around nine developmental stages of the newborn infant during skin-to-skin contact in the first hours after birth (17). It expands the understanding of each of these through the lens of the authors' clinical expertise and research experience from direct observation of newborn infant behaviour. Expertise comes from experience as well as analysis of hundreds of hours of videotapes of newborn infants' developing feeding behaviour in skin-to-skin contact

**Table 1** Shows the nine stages

Stages	Behaviours
1. Birth cry	Intense cry just after birth, transition to breathing air.
2. Relaxation stage	Infant rests. No activity of mouth, head, arms, legs or body.
3. Awakening stage	Infant begins to show signs of activity. Small thrusts of head: up, down, from side-to-side. Small movements of limbs and shoulders.
4. Active stage	Infant moves limbs and head, more determined movements. Rooting activity, 'pushing' with limbs without shifting body.
5. Resting stage*	Infant rests, with some activity, such as mouth activity, sucks on hand.
7. Familiarization	Infant has reached areola/nipple with mouth positioned to brush and lick areola/nipple.
8. Suckling stage	Infant has taken nipple in mouth and commences suckling.
9. Sleeping stage	Infant closes eyes and falls asleep.

\*The Resting Stage could be interspersed with all the stages.

(12,17,18,21,28–42). Additionally, the authors had the opportunity to improve skin-to-skin care after birth, both in vaginal and caesarean delivery at 10 hospitals in Egypt and the United States according to a 5-day video-ethnographic intervention. This methodology has five features: (i) a lecture educating staff (obstetricians, paediatricians, midwives, nurses and other staff at the delivery – and neonatal wards) about the theory behind the procedure of skin-to-skin care; (ii) practical application of the skin-to-skin procedure, with the American and Swedish team and staff working at the delivery ward together, to continue the educational process; (iii) videotaping the evolving process as the hospital staff work to implement the new procedures; (iv) conducting an interaction analysis workshop to review videotapes and discuss barriers and solutions; and (v) administering the continuing application of the procedure (12,28).

## RESULTS

### Before labour

Ideally, education related to the placement of the baby skin-to-skin with the mother should also have occurred as part of her antenatal care. When the mother and her companion (s) arrive at the birthing facility, a review of their understanding of the newborn infant's instinctive behaviours and the process of safe skin-to-skin immediately after birth should take place (Table 1). It has been our experience that parents are captivated by the newborn infant's instinctive behaviours and respond positively when they learn about the 9 stages and can identify the newborn infant's actions (32). Parents should also have an understanding about the importance of monitoring the newborn infant's position, breathing and safety. Providing the parents with a pamphlet or tear sheet or having a poster of the 9 instinctive stages somewhere in the birthing suite has been found to be helpful for the parents to follow the baby throughout the stages and recognise their newborn infant's ability. This also gives a clear role to the companion and provides both motivation to remain with the mother and newborn infant, and an additional opportunity for the baby's safety. Parents thereby have this additional knowledge and can prioritise focus on the newborn infant rather than distractive actions such as phone calls (43,44), posting on the Internet, talking to friends or family in the room, which are behaviours that are best postponed until after this unique time.

Staff have the opportunity to protect this important, small window of time for the parents to follow their newborn infant's behavioural development. This creates a shelter for the mother and baby's right to not be separated (45), to stay skin-to-skin after birth until the newborn infant has suckled and/or fallen asleep. Interruption of skin-to-skin contact during the first two hours reduces the infant's chances of early breastfeed (13,46).

Staff should also assure practical arrangements have been made for the experience of skin-to-skin. This includes ensuring that the mother's clothing has been arranged so it will be easy to remove and will allow access to the mother's chest immediately after the birth, being aware of any

intravenous lines in relation to the sleeves of the mother's gown, etc. Regardless of the birthing position, the mother needs a comfortable and supportive position during the first hours while skin-to-skin with the newborn infant. Hospital protocol should be reviewed to ensure it reflects best practice.

## THE 9 STAGES – A BREASTFEEDING START LED BY THE NEWBORN INFANT

### Stage 1 – The birth cry

This first stage is characterised by the initial birth cry, when the lungs expand for the first time as the newborn infant transitions to breathing, and other survival instincts (Fig. 1A). These behaviours could include the moro reflex, grimacing, coughing, lifting the full body from the mother's torso, abruptly opening the eyes and tension in the body. The baby's motions during the birth cry stage emanates from the drive to survive. During this extremely alert period, the newborn infant is able to make defensive movements with the hands to protect their airway, for example against a suction catheter or bulb.

The initial birth cry, and the subsequent crying during the first minutes after the birth, has the effect of expectorating the airways of the amniotic fluid (47). In addition, extremely high catecholamine levels at birth help in absorbing liquid from the airways (48). The American Association of Pediatrics (49) and Academy of Gynecologists and Obstetricians (ACOG) (50) have strong statements about the ability of healthy full-term babies to clear their own airways and successfully transition without the need of any form of suctioning. Research has shown that suctioning may disrupt the inborn sequential behaviours of the newborn infant (34). Interruption of the natural newborn behaviours immediately after birth could affect cell proliferation in the locus coeruleus in rats (51), with consequences of impaired locus coeruleus connected autism (52) and to stress-induced fear-circulatory disorders (53).

While transitioning the baby to the mother's chest, it is important to avoid compressing the baby's thorax, which could hamper breathing. Clinical practice should include gentle hands to hold the newborn infant in the drainage position (tilted with the head lower than the torso and the head slightly to the side) immediately after the birth, allowing the fluid to flow freely from the mouth and nose.

The comfortably positioned, semi-reclined mother receives the baby, to hold prone, skin-to-skin. The mother's semi-reclined position is conducive to the baby's breathing adaptation, in contrast to a horizontal position (54). If possible, the baby should be in a lengthwise position on the mother's body, with the head on the mother's chest, and above her breasts. The lengthwise mid-chest position also emphasises the importance of the upcoming stages. Positioning the baby's mouth close to the mother's nipple would put the focus on immediate breastfeeding, which the newborn infant is not ready to do now.

Standard practice includes drying the baby's head and body with a clean, dry cloth to help maintain body



**Figure 1** (A–I): Pictures of the 9 stages ©Healthy Children Project, Inc. Used with permission.

temperature. The baby's face should initially be turned to the side, which facilitates free airways and monitoring of the baby's breathing. After settling in the skin-to-skin position with the mother, the baby's body should be covered with a dry cloth, leaving the face uncovered.

During a caesarean surgery, the newborn infant might be positioned horizontally above the drape instead of vertically. In this case, the newborn infant is placed across the mother's breasts. This position requires additional attention from the staff to watch the baby's breathing and to ensure that the newborn infant does not push or jump-off of the mother.

Placement of the newborn infant skin-to-skin increases uterine contraction immediately after birth, increases the completeness of the delivered placenta and decreases uterine atony and excessive blood loss (3). Skin-to-skin also significantly decreases the duration of the third stage of labour (3).

Delayed cord clamping (>180 seconds after delivery) is recommended when compared to early cord clamping (>10 seconds after delivery). Delayed cord clamping reduced the risk of anaemia at 8 and 12 months in a group of high-risk children. In addition, there was a possible positive effect in infant's health and development (55,56). 'From a physiological point of view the appropriate time for umbilical cord clamping in healthy full-term infants is after

the infant has aerated its lungs, commenced breathing and the pulmonary blood flow has increased to sustain ventricular preload and cardiac output' (57). Deferred cord clamping is in line with recommendations from the World Health Organization (58).

Clinical practice shows that the umbilical cord should be left long, not clamped close to the newborn infant's belly, so that the newborn infant is not disturbed by the clamp, since clamps, and hard clamps especially, can be uncomfortable in a prone position. This can result in the babies lifting their body from the mother's chest, which reduces skin-to-skin contact and thus compromises the newborn infant's temperature. It can also cause a cry of discomfort.

Staff should create a teaching situation at their routine frequent observations of the baby during skin-to-skin contact, involving the parents in unobtrusive, gentle, close observation, 'helping them understand and avoid potentially hazardous positions' (59) during the first few hours after the birth. In addition, the nine stages offer an opportunity for staff to discuss the progress of the newborn infant and to ensure continued observation by the parents or attendants. The focus on following the newborn infant's progress through the nine instinctive stages results in a close examination of the baby by the mother's companion(s) and staff throughout the vulnerable first two hours. This may help to illuminate the capabilities and limitations of a

specific baby, and thereby increase the safety during this important time. The safety of the newborn infant and mother is always the highest priority (Table 2).

We have seen that when an unmedicated mother receives her newborn infant into her arms immediately after birth, she grasps the baby with confidence. She will often hold the baby gently around the torso, look at the baby and then bring the baby to her heart, allowing the newborn infant to lie quietly skin-to-skin with her, as both relax. When the mother has the newborn infant immediately after birth on her naked chest, it has a profound impact on her. A systematic review of mother's experiences of skin-to-skin contact includes overwhelming feelings of love, a natural experience that taught them how to be a mother, improved self-esteem, and a way of knowing and understanding the infant (60). We have noticed that this simple act of the staff handing over the newborn infant to the mother supports early parental confidence.

**Table 2** Safe skin-to-skin care

Safe Interactive Skin-to-Skin Contact in the First Hour After Birth

1. Make sure that the mother is in a comfortable semi-reclined position with support under her arms.
2. After drying the newborn infant, lift the newborn infant gently to avoid compression of the thorax when placing the baby skin-to-skin. Put the baby prone, in a lengthwise position with the head on the mother's chest above the breast.
3. Cover the baby with a dry blanket/towel. Leave the face visible.
4. Make sure that the nose and mouth are not enveloped by the mother's breast or body or obscured by the blanket. Initially, the baby's head should be turned to the side.
5. The newborn infant must have the opportunity to use its reflexes to lift the head so the nose and mouth can be free. This is of special concern if the mother has large and/or very soft breasts.
6. The nipple must be accessible to the newborn infant. For some mothers, this may require positioning a towel or pillow under or on the side of the mother's breast.
7. Show the parents how to support the breast to secure free airways especially during the time the baby starts searching for the breast. Verify understanding.
8. Remind the parents to focus on the newborn infant and follow the newborn infant's early behavior, making sure that the parents follow the 9 stages. The other parent should be observant, not distracted by mobile phones, etc., during skin-to-skin.
9. Extra attention may be required if the mother is affected by sedation after childbirth as well as during possibly postpartum suturing. The other parent should be aware of the situation and watch for the safety of the infant.
10. Labor medications can affect the newborn infant, and hamper reflexes. The medications may impair the newborn infant's reflexes enough to prevent the ability to lift the head to protect itself from suffocation. Babies affected by labor medications must be constantly monitored.

*Adapted from* The Swedish-American Team for Baby Adapted Care of Health Infants in the First Hours after Birth, Widström, Svensson and Brimdyr, for Kom Ombo Hospital, Egypt, 2007 and from [www.lof.se/patientsakerhet](http://www.lof.se/patientsakerhet).

If the mother has received analgesics during labour, special care must be taken to watch for free airways. For example, pethidine can cross the placenta and affect the newborn infant's breastfeeding behaviour negatively (33) and may specifically hamper the newborn infant's ability to lift the head (38) as well as infant's temperature and crying (39). Fentanyl given in the epidural space passes rapidly into maternal blood. A placental transfer of 90% has been measured by Moises (61). Fentanyl has been measured in the newborn infant's urine for at least 24 hours postpartum (62). Fentanyl exposure can depress the newborn infant's behaviour during the first hours after birth, especially the suckling behaviour in a dose-dependent manner (29).

### Stage 2 – Relaxation

During the relaxation stage (Fig. 1B), the newborn infant is still and quiet, making no movements (Fig. 1B). Clinical experience shows it is not possible to elicit a rooting reflex during the relaxation stage. The baby's sensory system seems to be depressed. From an evolutionary perspective, this silent and motionless period may be a way to hide from predators during a vulnerable time (17). When lying quietly on the mother's chest, the baby can hear the mother's heartbeat. This familiar sound from in utero seems to comfort the newborn infant after the rapid transition to extra-uterine life.

It is suggested that the pressure on the head through the birth canal is the cause of extremely high catecholamine level after birth, a level 20 times higher than that of a resting adult (63). This high catecholamine concentration might partly be the cause of the higher pain threshold in the baby close to birth (64) and be a mirror of nature's way to relieve pain in the baby when passing through the birth canal. Consequently, the baby's temporarily impaired sensation at birth causes the relaxation stage – the baby has decreased sensitivity to the surroundings.

We have noticed that staff members exhibit concerns about the newborn infant's silent and relaxed state during this time. This stems from a lack of understanding of the baby's stage, often resulting in rubbing or massaging the newborn infant in a disruptive and vigorous manner, repositioning or lifting the baby from the mother. If, during the relaxation stage, the newborn infant is disturbed by the actions of the staff, the baby will react with crying, grimacing and protective reflexes.

If separated from their mother, babies exhibit a distinct separation distress call. This separation distress vocalisation in mammalian species that stops at reunion may be nature's way of keeping newborn infants warm with maternal body temperature. The baby's call at separation is thus a survival mechanism (65,66).

Mammalian species, including human mothers, react to these separation distress calls and intuitively attempt to retrieve the newborn infant. In hospital settings, where the mother has limited control over her environment, we have noticed her straining to look at the infant who has been taken away and asking others about the state of the infant. (Fig. 2)

It is possible to conduct the assessment of the APGAR score, as well as any other necessary assessments, on a healthy full-term newborn infant without disturbing the infant, allowing skin-to-skin to continue uninterrupted. It is more effective and advantageous to assess the newborn infant when skin-to-skin with mother since babies are less likely to cry; they are more likely to remain warm and not waste energy.

If administration of vitamin K shot is a routine, this should occur soon after birth while the catecholamine levels are highest (10), preferably with the newborn infant skin-to-skin with the mother, as skin-to-skin contact has been shown to lower the baby's reaction to pain in the postpartum (67).

### Stage 3 – Awakening

The awakening stage (Fig. 1C) is a transition from the relaxation stage to the activity stage. The baby makes small motions. Small movements of the head, face and shoulders gently ripple down through the arms to the fingers. The baby makes small mouth movements. They will gradually open their eyes during this stage, blinking repeatedly until the eyes are stable and focused (Fig. 1C).

### Stage 4 – Activity

During the activity stage (Fig. 1D), the baby exhibits a greater range of motion throughout the head, body, arms and hands. The limbs move with greater determination; the baby may root and lift the head from the mother's chest. The fingers often begin the stage as fisted but may expand to massage, transfer tastes with a hand-to-nipple-to-mouth movement, catch the nipple and explore the mother's chest (17). Rooting also becomes more obvious during this stage (34).

The successive protrusion of the tongue continues throughout this stage. At the beginning of this stage, the newborn infant may have only moved the tongue within the mouth. During the activity stage, the baby will bring the tongue to the edge of the lips, then protrude beyond the lips, then protrude repeatedly beyond the lips (40). These tongue exercises, which may pave the way for later tongue behaviours, specifically suckling, can be affected by medications, such as pethidine (33).

During pregnancy, the nipple has become more pigmented (68) and is easy for the newborn infant to discover (Fig. 1C). We have observed that soon after birth, the areola expands and takes a bulb-like shape (Fig. 1G). The Montgomery glands also become more pronounced (Fig. 1G). The scent of areolar secretions has been linked to behavioural responses, such as head turning (69) and directional crawling in newborn infants (70). This release of the breast odour by the Montgomery glands is known to help the newborn infant find the nipple (17,27,69). The newborn infant recognises the scent of the mother's breast from the amniotic fluid (71), touches the breast and transmits the taste of the breast to the mouth (hand-to-breast-mouth movements) (17). This stimulates rooting and crawling movements in the newborn infant to reach the nipple. The connection between the taste of amniotic fluid

and the scent of the breast from the Montgomery glands highlights a biological survival mechanism – a pathway of flavour with lifelong consequences. (Fig. 3) ([17,71–80]) When knowing of this sensitive odour-dependent mechanism, it might be wise not to interfere with unfamiliar hands.

The infant has learned to recognise the mother's voice during intrauterine life (81). The infant's learning and memory skills are quite sophisticated; the foetus can learn to recognise a vowel from the surrounding language and is some days after birth able to discriminate this vowel from a vowel from a foreign language (82).

After the newborn infant has located the nipple by sight, the mother's voice will attract the baby's attention to her face. Arousal transfer between mother and 4- to 6-month-old babies is likely to happen through pupillary contagion (respond to pupil size with changes in one's own) is fundamental for social and emotional development (83). Interestingly, it is described in a paediatric essay that 'immediately after birth the newborn's eyes become wide open, usually with large pupils' (63). When skin-to-skin with the mother after birth and free to safely move, the newborn infant searches for eye contact with the mother around half an hour after birth (17) (Fig. 1E). We suggest that infant-maternal bonding through pupillary contagion may start early after birth as our clinical experience is that mothers often recall the first moment of eye-to-eye contact as unforgettable. The complex experiences of the newborn infant during the first hour encompass more than simply a journey to the breast; the opportunity for eye contact emphasises the importance of parents and staff valuing instinctive behaviour during this time, and the avoidance of interruptions.

### Stage 5 – Resting

The resting stage (Fig. 1E) is intertwined with all of the other stages. A baby may stop or start during any of the stages to rest, and then continue with that same stage, or move on to the next (12,84). The baby could be lying still sucking on fingers or just gazing at the nipple. The eyes may be open or closed.

It is important to value the resting stage and not worry that the baby is done and has not been successful with the process of the first hour. We have seen this stage being misinterpreted by parents or staff, resulting in the newborn infant being removed from skin-to-skin with the mother in favour of other routine care. It is vital to allow the newborn infant to take these pauses throughout the first hour or so without being interrupted or separated, remembering that this stage is naturally interspersed throughout other stages. Research about adult 'awake rest' points to the importance of this time to the consolidation of memories, and it's contribution to learning (85). This could be applicable to the newborn infant's rest during the first hour as well (41). Separation causes a critical break of the stages (31). If the newborn infant is separated from the mother, even if the infant is returned, the newborn infant might need to begin again at the first stage – crying and relaxing before



**Figure 2** Caesarean mother looking towards her baby ©Healthy Children Project, Inc. Used with permission.

beginning to progress through the stages again, which might take some time. The likelihood increases that the newborn infant may not be able to go through all the stages before needing to sleep, compromising the possibility to suckle, Stage 8.

#### Stage 6 – Crawling

The crawling stage (Fig. 1F) could include crawling, leaping, sliding, bulldozing or many other interesting names of ways to move from the position between the breasts to a position very close to the nipple. Sometimes this process is so subtle that parents and staff are surprised to notice that the baby has made its way over to the breast. Other times, the baby may make strong and overt motions, collecting the attention of everyone in the room. The evolutionary purpose behind the newborn infant's innate so-called stepping reflex becomes clear as the newborn infant crawls to the mother's breast. The movements of these steps of the feet over the mother's uterus may contribute to the contractions of the uterus, and the decreased time to expel the placenta (2,3) and decreased blood loss (3) during skin-to-skin.

During this process, it is important to protect the newborn infant's effort to reach the breast, with the intention not to lift or turn the baby's body. Placing a towel or pillow under the mother's arms is important during this stage, since the newborn infant will be travelling over to one side of the chest and therefore (in many settings) close to the edge of the bed (18). This support can help the newborn infant to reach the nipple without becoming exhausted by unnecessary sliding in the wrong direction. It is also important for parents to be aware of the travelling process, since they may try to reorient the baby back to the middle of the chest, to keep the newborn infant from the edges of the bed. Support under the mother's arms may also allow the nipple to remain in an area easy for the baby to find and grasp. A mother might use her own palm to arrange the breast as well. A baby needs to use the feet to achieve crawling, and sometimes the feet are in a less than ideal position. They may be off the side of the mother's body or

pushing in the air. In those cases, it may be helpful if the mother puts her hand under the newborn infant's foot to give the baby something to push against in order to move towards the breast.


#### Stage 7 – Familiarization

Since the baby is prone on a semi-reclined mother, the alert newborn infant has the control over their experience, rather than being placed in a dependent position by the mother (Fig. 1G). To reach the breast, it must be possible for the baby to manoeuvre into an appropriate position. When approaching the breast the newborn infant performs specific soliciting calls to mother – a short clinging call that usually results in a gentle response from the mother. The frequency of these sounds increases as the newborn infant gets closer to the mother's nipple (17). The odours from the mother's breast are likely to be inducing this response. Parents respond to the newborn infant's soliciting (42).

During the familiarization stage (Fig. 1G), the baby becomes familiar with the breast by licking the nipple and areola. This period could last 20 minutes or more. The newborn infant massages the breast, which increases the mother's oxytocin levels (35) and shapes the nipple by licking. During this stage, it is evident that the baby is smelling and tasting, and previous actions become more vigorous and more coordinated. Therefore, it is important not to interfere or introduce odours from unfamiliar hands.

The baby continues with tongue activity during this stage, now more overtly related to eventual breastfeeding, by licking above and below the nipple. The baby may make noises with the mouth and lips, like smacking sounds, during this stage. The breast and nipple are shaped by the massaging and licking actions of the newborn infant. The newborn infant is preparing the tongue, breast and nipple for the moment of attachment and suckling.

The actions of the tongue inform the staff of the newborn infant's coordination of the tongue with the rooting reflex, and the ability to move the tongue to the bottom of the mouth, curved and thin. The newborn infant needs to practice this coordination of the rooting-tongue reflex (36).



**Figure 3** Pathway of flavours.

Staff and parents should allow the baby the time needed to practice this coordination during this stage; the newborn infant is perfecting many important oral-motor functions. This learning is vital as the newborn infant initiates the suckling process.

There is often a resting stage between the familiarization stage and the suckling stage, which unfortunately can make parents and staff prone to (so-called) help the newborn infant to the breast.

It is also common for the baby to attach, suck once or twice, and then disengage. The newborn infant will be conducting a normal step of the familiarization Stage, but it may look to staff like the newborn infant is unable to attach. The baby must thus be allowed to do these moves to adjust into an instinctive position. This is conducive to the newborn infant's chin making an initial contact with the mother's breast as the baby endeavours to catch the nipple. 'Chin-first contact' is associated with sustained deep, rhythmical suckling (86).

In a study of babies who had later been diagnosed with significant latch problems, the majority of mothers reported that the newborn infant had been forced to latch to his mother's breast. According to the mothers, the babies screamed, acted in a panicky way, exhibited avoidance behaviours and had other strong reactions against the forceful treatment (87). It has also been shown that mothers who had this type of so-called help have a more negative experience of the first breastfeeding (88) and breastfeed for a shorter time (88,89).

If infants with an aversive behaviour are allowed to peacefully go through the stages in skin-to-skin contact at a later time, they may successfully reach the nipple, attach to

the nipple by themselves and start suckling. This could happen even weeks after birth if not possible earlier (87). This is a promising way to calmly solve breastfeeding problems, especially since a new study shows that breastfeeding is associated with decreased childhood maltreatment (90).

### Stage 8 – Suckling

The newborn infant attaches to the nipple during this stage and successfully breastfeeds (Fig. 1H). When babies self-attach, they are positioning the wide-open mouth appropriately on the areola and nipple, protecting against sore nipples. It is interesting to note that the hands, which have been so busy, often stop moving once suckling begins (35); the eyes, which have been looking at the breast, the mother and the room, often become focused after attaching.

During this first hour, when the unmedicated baby self-attaches, it is a perfect first breastfeeding, although the infant will continue to readjust until satisfied with the latch. The newborn infant does not need *help* to adjust the latch. Babies who self-attach during the first hour after birth have few problems with breastfeeding, latch and milk transfer (13). Skin-to-skin in the first hour strengthens the mother's self-confidence, including decreasing the concerns about having enough milk (4). When babies are placed skin-to-skin with the mother, they have more optimal blood glucose levels. Both skin-to-skin and the suckling contribute to this effect (91). Thus, this reduces the risk of supplementation.

Medicated babies can successfully go through the nine stages and self-attach. However, there is increasing evidence concerning the negative consequences of certain medications such as fentanyl and oxytocin, on success with

breastfeeding (29). Parents and staff must take into account the consequences when considering amount, timing and choice of specific labour medications.

### Stage 9 – Sleeping

Towards the end of suckling, about an hour and a half after birth, the newborn infant becomes drowsy and falls asleep (Fig. 11). The oxytocin, released in both mother and infant by suckling, triggers the release of gastrointestinal (GI) hormones, including cholecystokinin (CCK) and gastrin. The high level of CCK in both mother and newborn infant will cause a relaxing and satisfying postprandial sleep (37). The GI activity will also improve maternal and infant nutritional absorption. The advantages, and the *feel-good* effect, will continue at each breastfeeding session.

If the mother and newborn infant were unable to experience these first hours together, or if the newborn infant did not self-attach before falling asleep, the dyad should have the opportunity to stay skin-to-skin as much as possible. If separation occurs, hand expression of milk within the first hour after birth enhances milk production (92). The partner can also spend time skin-to-skin with a new baby if separation from the mother is necessary (42).

When the baby is alert and shows signs of readiness to feed, the processes of finding the mother's breast when skin-to-skin usually happen more quickly in the postpartum period than soon after birth. One reason for this is that the initial stages of a birth cry, relaxation and awakening are associated with the situation and hormonal status unique to the time immediately after birth. Later after birth, the baby has more control over his body and movements.

Understanding the stages, and the breastfeeding behaviours of the newborn infant, is indeed, reassuring to the parents, even in the postpartum ward.

There have been discussions about a possible relationship between skin-to-skin contact and Sudden Unexpected Postnatal Collapse (SUPC) (43). The background to SUPC is multifactorial. However, one of the common causes of the collapse seems to be respiratory distress (44,93–95). The possibility of labour medications affecting the newborn infant adversely is not highlighted in the Sudden Unexpected Postnatal Collapse (SUPC) literature. (43,96,97). The majority of the cases occur in early skin-to-skin care, without observance of the child, or when sleeping with accidentally covered respiratory tract (98,99). A review of Sudden Unexpected Early Neonatal Deaths (SUENDs) in Sweden shows that a higher incidence of SUENDs was reported before the practice of early skin-to-skin contact was introduced (100,101). Staff need to be aware of the importance of safe skin-to-skin, including the close observation of the newborn infant's airways (Table 2).

### Possible long-term benefits of skin-to-skin contact

Immediate skin-to-skin contact provides the initial colonisation of the baby's microbiome outside of the mother, a swarming of the mother's skin bacteria. The microbial colonisation of the infant begins before birth and continues through the birth canal. This is one of the reasons the

newborn infant should not be washed during this time. Colonisation after caesarean surgery does not match vaginal colonisation (102); skin-to-skin may be of extra importance in these cases. As the hour progresses, the first tastes of colostrum will provide vital sustenance to the infant's developing gut microbiota, which has been implicated in the expression of genes (103). Evolving research on epigenetics and the microbiome highlights the importance of the optimal microbiome (104), enhanced by breastfeeding, which has been implicated in long-term health, including decreased obesity and metabolic diseases (103).

In a longitudinal study, it was found that the mother's breast temperature increases when mother and newborn infant are in skin-to-skin contact, resulting in an increase of newborn infant's foot temperature (8). The warmer foot temperature is an indication of the decrease of the negative effect of the 'stress of being born' that is associated with skin-to-skin. Those mothers who hold their babies skin-to-skin after birth were rated as 'less rough' when latching and stimulating their babies during breastfeeding at day 4 postpartum (105). Skin-to-skin was also linked to improved mother/infant mutuality one year later. Skin-to-skin after birth also positively influenced the infant's self-regulation at one year (106). Self-regulation is a part of the concept of self-control.

Interestingly, Moffitt et al. showed that if a child has good self-control at the age of 4, this has consequences into adulthood, in terms of increased education and income and decreased drug addiction and criminal behaviours as measured at 30 years of age, compared to those with low self-control at four years of age, among a cohort of 1,000 children (107). Thus, skin-to-skin contact after birth may contribute to positive long-term consequences.

From the beginning, biological processes during the first hour after birth ensure survival of the mother and infant. This sensitive period has an emotional consequence on the mother's understanding of the newborn infant, enhancing bonding. Humans are resilient, and there are opportunities to bond and breastfeed even when the experience immediately after the birth is less than optimal. For example, even days and months after the birth, the infant will exhibit these same instinctive breastfeeding behaviours, with the potential to overcome early breastfeeding problems. But this sensitive period immediately after birth must be recognised and valued, both by parents and staff, to provide the opportunity for this unique experience. It is vital not to interrupt these natural behaviours, as they form nature's basis for attachment, and support the mother's confidence in her infant's inborn capability. This may have significant consequences for the parent's understanding of the baby throughout childhood, protecting the child from parental *roughness* and laying a foundation for the child's self-regulation and self-control.

### CONCLUSION

A sensitive period of the newborn infant coincides with mother's sensitive period during the first hour after birth. This is a unique situation for both, individually and in

relation to each other, and must be protected by evidence-based routines of hospital staff. Clinical expertise, together with systematic evidence, combines to enhance the understanding of the best practice of skin-to-skin care in the first hours after birth. Hospital guidelines must support and enhance this important time.

## FUNDING

The study did not receive any specific funding.

## CONFLICTS OF INTEREST

The authors have no conflicts of interest to declare.

## References

- Moore ER, Bergman N, Anderson GC, Medley N. Early skin-to-skin contact for mothers and their healthy newborn infants. *Cochrane Database Syst Rev* 2016; 11: CD003519.
- Marín Gabriel MA, Llana Martín I, López Escobar A, Fernández Villalba E, Romero Blanco I, Touza Pol P. Randomized controlled trial of early skin-to-skin contact: effects on the mother and the newborn. *Acta Paediatr Oslo Nor* 1992; 2010: 1630–4.
- Essa RM, Abdel Aziz Ismail NI. Effect of early maternal/newborn skin-to-skin contact after birth on the duration of third stage of labor and initiation of breastfeeding. *J Nurs Educ Pract* 2015; 5. Available from: <http://www.sciedu.ca/journal/index.php/jnep/article/view/5698>
- Aghdas K, Talat K, Sepideh B. Effect of immediate and continuous mother-infant skin-to-skin contact on breastfeeding self-efficacy of primiparous women: a randomised control trial. *Women Birth J Aust Coll Midwives* 2014; 27: 37–40.
- Handlin L, Jonas W, Petersson M, Ejdebäck M, Ransjö-Arvidson A-B, Nissen E, et al. Effects of sucking and skin-to-skin contact on maternal ACTH and cortisol levels during the second day postpartum-influence of epidural analgesia and oxytocin in the perinatal period. *Breastfeed Med Off J Acad Breastfeed Med* 2009; 4: 207–20.
- Nissen E, Lilja G, Widström AM, Uvnäs-Moberg K. Elevation of oxytocin levels early post partum in women. *Acta Obstet Gynecol Scand* 1995; 74: 530–3.
- Winberg J. Mother and newborn baby: mutual regulation of physiology and behavior—a selective review. *Dev Psychobiol* 2005; 47: 217–29.
- Bystrova K, Widström AM, Matthiesen AS, Ransjö-Arvidson AB, Welles-Nyström B, Wassberg C, et al. Skin-to-skin contact may reduce negative consequences of “the stress of being born”: a study on temperature in newborn infants, subjected to different ward routines in St. Petersburg. *Acta Paediatr Oslo Nor* 1992; 2003: 320–6.
- Beiranvand S, Valizadeh F, Hosseinabadi R, Pourmia Y. The effects of skin-to-skin contact on temperature and breastfeeding successfulness in full-term newborns after cesarean delivery. *Int J Pediatr* 2014; 2014: 1–7.
- Hägnevik K, Faxelius G, Irestedt L, Lagercrantz H, Lundell B, Persson B. Catecholamine surge and metabolic adaptation in the newborn after vaginal delivery and caesarean section. *Acta Paediatr Scand* 1984; 73: 602–9.
- Bramson L, Lee JW, Moore E, Montgomery S, Neish C, Bahjri K, et al. Effect of early skin-to-skin mother–infant contact during the first 3 hours following birth on exclusive breastfeeding during the maternity hospital stay. *J Hum Lact Off J Int Lact Consult Assoc* 2010; 26: 130–7.
- Crenshaw JT, Cadwell K, Brimdyr K, Widström A-M, Svensson K, Champion JD, et al. Use of a video-ethnographic intervention (PRECESS Immersion Method) to improve skin-to-skin care and breastfeeding rates. *Breastfeed Med Off J Acad Breastfeed Med* 2012; 7: 69–78.
- Righard L, Alade MO. Effect of delivery room routines on success of first breast-feed. *Lancet* 1990; 336: 1105–7.
- World Health Organization. Implementation guidance: protecting, promoting and supporting breastfeeding in facilities providing maternity and newborn services – the revised Baby-friendly Hospital Initiative. Geneva; 2018. Report No.: CC BY-NC-SA 3.0 IGO.
- Klaus MH, Jerauld R, Kreger NC, McAlpine W, Steffa M, Kennell JH. Maternal attachment: importance of the first postpartum days. *N Engl J Med* 1972; 286: 460–3.
- Kennell JH, Trause MA, Klaus MH. Evidence for a sensitive period in the human mother. In: Porter R, O'Connor M, editos. *Novartis Foundation Symposia*. Chichester, UK: John Wiley & Sons, Ltd, 2008: 87–101. Available at: <https://doi.org/doi.wiley.com/10.1002/9780470720158.ch6>
- Widström A-M, Lilja G, Aaltomaa-Michalias P, Dahllöf A, Lintula M, Nissen E. Newborn behaviour to locate the breast when skin-to-skin: a possible method for enabling early self-regulation. *Acta Paediatr Oslo Nor* 1992; 2011: 79–85.
- Brimdyr K, Wiström AM, Svensson K. *Skin to Skin in the First Hour after Birth: Practical advice for staff after vaginal and cesarean birth [DVD]*. Sandwich, MA, USA: Healthy Children Project, Inc., 2011.
- Keverne EB, Levy F, Poindron P, Lindsay DR. Vaginal stimulation: an important determinant of maternal bonding in sheep. *Science* 1985; 219: 81–3.
- Theodosis DT, Chapman DB, Montagnese C, Poulain DA, Morris JF. Structural plasticity in the hypothalamic supraoptic nucleus at lactation affects oxytocin-, but not vasopressin-secreting neurones. *Neuroscience* 1986; 17: 661–78.
- Widström AM, Wahlberg V, Matthiesen AS, Eneroth P, Uvnäs-Moberg K, Werner S, et al. Short-term effects of early suckling and touch of the nipple on maternal behaviour. *Early Hum Dev* 1990; 21: 153–63.
- Fleming AS, Corter C, Franks P, Surbey M, Schneider B, Steiner M. Postpartum factors related to mother's attraction to newborn infant odors. *Dev Psychobiol* 1993; 26: 115–32.
- Corona R, Lévy F. Chemical olfactory signals and parenthood in mammals. *Horm Behav* 2015; 68: 77–90.
- Nelson EE, Panksepp J. Brain substrates of infant-mother attachment: contributions of opioids, oxytocin, and norepinephrine. *Neurosci Biobehav Rev* 1998; 22: 437–52.
- Raineki C, Pickenhagen A, Roth TL, Babstock DM, McLean JH, Harley CW, et al. The neurobiology of infant maternal odor learning. *Braz J Med Biol Res Bras Pesqui Medicas E Biol* 2010; 43: 914–9.
- Bartocci M, Winberg J, Ruggiero C, Bergqvist LL, Serra G, Lagercrantz H. Activation of olfactory cortex in newborn infants after odor stimulation: a functional near-infrared spectroscopy study. *Pediatr Res* 2000; 48: 18–23.
- Varendi H, Porter RH, Winberg J. Does the newborn baby find the nipple by smell? *Lancet Lond Engl* 1994; 344: 989–90.
- Brimdyr K, Widström A-M, Cadwell K, Svensson K, Turner-Maffei C. A realistic evaluation of two training programs on implementing skin-to-skin as a standard of care. *J Perinat Educ* 2012; 21: 149–57.
- Brimdyr K, Cadwell K, Widström A-M, Svensson K, Neumann M, Hart EA, et al. The association between common labor

- drugs and suckling when skin-to-skin during the first hour after birth. *Birth Berkeley Calif* 2015; 42: 319–28.
30. Cadwell K, Brimdyr K, Phillips R. Mapping, measuring, and analyzing the process of skin-to-skin contact and early breastfeeding in the first hour after birth. *Breastfeed Med Off J Acad Breastfeed Med* 2018; 13(7): 485–492.
  31. Brimdyr K, Cadwell K, Stevens J, Takahashi Y. An implementation algorithm to improve skin-to-skin practice in the first hour after birth. *Matern Child Nutr* 2018; 14: e12571.
  32. Brimdyr K, Widström A-M. *The Magical Hour: Holding your baby skin to skin for the first hour after birth [DVD]*. Sandwich, MA, USA: Healthy Children Project, Inc., 2012.
  33. Nissen E, Lilja G, Matthiesen AS, Ransjö-Arvidsson AB, Uvnäs-Moberg K, Widström AM. Effects of maternal pethidine on infants' developing breast feeding behaviour. *Acta Paediatr Oslo Nor* 1992; 1995: 140–5.
  34. Widström AM, Ransjö-Arvidsson AB, Christensson K, Matthiesen AS, Winberg J, Uvnäs-Moberg K. Gastric suction in healthy newborn infants. Effects on circulation and developing feeding behaviour. *Acta Paediatr Scand* 1987; 76: 566–72.
  35. Matthiesen AS, Ransjö-Arvidsson AB, Nissen E, Uvnäs-Moberg K. Postpartum maternal oxytocin release by newborns: effects of infant hand massage and sucking. *Birth Berkeley Calif* 2001; 28: 13–9.
  36. Widström AM, Thingström-Paulsson J. The position of the tongue during rooting reflexes elicited in newborn infants before the first suckle. *Acta Paediatr Oslo Nor* 1992; 1993: 281–3.
  37. Uvnäs-Moberg K, Widström AM, Marchini G, Winberg J. Release of GI hormones in mother and infant by sensory stimulation. *Acta Paediatr Scand* 1987; 76: 851–60.
  38. Nissen E, Widström AM, Lilja G, Matthiesen AS, Uvnäs-Moberg K, Jacobsson G, et al. Effects of routinely given pethidine during labour on infants' developing breastfeeding behaviour. Effects of dose-delivery time interval and various concentrations of pethidine/norpethidine in cord plasma. *Acta Paediatr Oslo Nor* 1992 1997; 86: 201–8.
  39. Ransjö-Arvidsson AB, Matthiesen AS, Lilja G, Nissen E, Widström AM, Uvnäs-Moberg K. Maternal analgesia during labor disturbs newborn behavior: effects on breastfeeding, temperature, and crying. *Birth Berkeley Calif* 2001; 28: 5–12.
  40. Brimdyr K, Widström AM, Cadwell K, Svensson K. Analysis of newborn tongue behavior as related to intrapartum epidural fentanyl exposure. 16th ISRHML Conference Breastfeeding and the Use of Human Milk, Science and Practice; 2012; Trieste, Italy.
  41. Cadwell K, Brimdyr K. Intrapartum administration of synthetic oxytocin and downstream effects on breastfeeding: elucidating physiologic pathways. *Ann Nurs Res Pr* 2017; 2: 1024.
  42. Velandia M, Matthiesen A-S, Uvnäs-Moberg K, Nissen E. Onset of vocal interaction between parents and newborns in skin-to-skin contact immediately after elective cesarean section. *Birth Berkeley Calif* 2010; 37: 192–201.
  43. Herlenius E, Kuhn P. Sudden unexpected postnatal collapse of newborn infants: a review of cases, definitions, risks, and preventive measures. *Transl Stroke Res* 2013; 4: 236–47.
  44. Feldman-Winter L, Goldsmith JP, Committee on fetus and newborn, Task force on sudden infant death syndrome. Safe sleep and skin-to-skin care in the neonatal period for healthy term newborns. *Pediatrics* 2016; 138: e20161889–e20161889.
  45. Convention on the Rights of the Child [Internet]. UN General Assembly; 1989 Nov. (United Nations, Treaty Series, vol. 1577, p. 3). Available at: <http://www.refworld.org/docid/3ae6b38f0.html>
  46. Robiquet P, Zamiara P-E, Rakza T, Deruelle P, Mestdagh B, Blondel G, et al. Observation of skin-to-skin contact and analysis of factors linked to failure to breastfeed within 2 hours after birth. *Breastfeed Med* 2016; 11: 126–32.
  47. Faxelius G, Hågnevik K, Lagercrantz H, Lundell B, Irestedt L. Catecholamine surge and lung function after delivery. *Arch Dis Child* 1983; 58: 262–6.
  48. Lagercrantz H. The stress of being born. *Sci Am* 1986; 254: 100–7.
  49. Wyckoff MH, Aziz K, Escobedo MB, Kapadia VS, Kattwinkel J, Perlman JM, et al. Part 13: Neonatal Resuscitation: 2015 American Heart Association Guidelines Update for Cardiopulmonary Resuscitation and Emergency Cardiovascular Care (Reprint). *Pediatrics* 2015; 136: S196–218.
  50. Committee Opinion No. 689. Delivery of a Newborn With Meconium-Stained Amniotic Fluid [Internet]. American College of Obstetricians and Gynecologists; 2017 p. 129:e33–4. Available at: <https://www.acog.org/Clinical-Guidance-and-Publications/Committee-Opinions/Committee-on-Obstetric-Practice/Delivery-of-a-Newborn-With-Meconium-Stained-Amniotic-Fluid>.
  51. Duarte ECW, de Azevedo MS, Pereira GAM, Fernandes MD, Lucion AB. Intervention with the mother–infant relationship reduces cell proliferation in the Locus Coeruleus of female rat pups. *Behav Neurosci* 2017; 131: 83–91.
  52. Mehler MF, Purpura DP. Autism, fever, epigenetics and the locus coeruleus. *Brain Res Rev* 2009; 59: 388–92.
  53. Bracha HS, Garcia-Rill E, Mrak RE, Skinner R. Postmortem locus coeruleus neuron count in three american veterans with probable or possible war-related PTSD. *J Neuropsychiatry Clin Neurosci* 2005; 17: 503–9.
  54. Erlandsson K, Christensson K, Dsilna A, Jnsson B. Do caregiving models after cesarean birth influence the infants' breathing adaptation and crying? A pilot study. *J Child Young Peoples Nurs* 2008; 2: 7–12.
  55. Andersson O, Hellström-Westas L, Andersson D, Domellöf M. Effect of delayed versus early umbilical cord clamping on neonatal outcomes and iron status at 4 months: a randomised controlled trial. *BMJ* 2011; 343: d7157.
  56. Kc A, Rana N, Målvqvist M, Jarawka Ranneberg L, Subedi K, Andersson O. Effects of delayed umbilical cord clamping vs early clamping on anemia in infants at 8 and 12 months: a randomized clinical trial. *JAMA Pediatr* 2017; 171: 264.
  57. Kluckow M, Hooper SB. Using physiology to guide time to cord clamping. *Semin Fetal Neonatal Med* 2015; 20: 225–31.
  58. Guideline: Delayed Umbilical Cord Clamping for Improved Maternal and Infant Health and Nutrition Outcomes [Internet]. Geneva: World Health Organization; 2014. (WHO Guidelines Approved by the Guidelines Review Committee). Available at: <http://www.ncbi.nlm.nih.gov/books/NBK310511/>
  59. Fleming PJ. Unexpected collapse of apparently healthy newborn infants: the benefits and potential risks of skin-to-skin contact. *Arch Dis Child - Fetal Neonatal Ed* 2012; 97: F2–3.
  60. Anderzén-Carlsson A, Lamy ZC, Eriksson M. Parental experiences of providing skin-to-skin care to their newborn infant—Part 1: A qualitative systematic review. *Int J Qual Stud Health Well-Being* 2014; 9: 24906.
  61. Moisés ECD, de Barros Duarte L, de Carvalho Cavalli R, Lanchote VL, Duarte G, da Cunha SP. Pharmacokinetics and transplacental distribution of fentanyl in epidural anesthesia for normal pregnant women. *Eur J Clin Pharmacol* 2005; 61: 517–22.
  62. Moore A, el-Bahrawy A, Hatzakorzian R, Li-Pi-Shan W. Maternal epidural fentanyl administered for labor analgesia is

- found in neonatal urine 24 hours after birth. *Breastfeed Med* 2016; 11: 40–1.
63. Lagercrantz H. The good stress of being born. *Acta Paediatr* 2016; 105: 1413–6.
  64. Bergqvist LL, Katz-Salamon M, Hertegård S, Anand KJS, Lagercrantz H. Mode of delivery modulates physiological and behavioral responses to neonatal pain. *J Perinatol* 2009; 29: 44–50.
  65. Christensson K, Cabrera T, Christensson E, Uvnäs-Moberg K, Winberg J. Separation distress call in the human neonate in the absence of maternal body contact. *Acta Paediatr Oslo Nor* 1992; 1995: 468–73.
  66. Michelsson K, Christensson K, Rothgänger H, Winberg J. Crying in separated and non-separated newborns: sound spectrographic analysis. *Acta Paediatr Oslo Nor* 1992; 1996: 471–5.
  67. Olsson E, Ahlsén G, Eriksson M. Skin-to-skin contact reduces near-infrared spectroscopy pain responses in premature infants during blood sampling. *Acta Paediatr* 2016; 105: 376–80.
  68. Stone K, Wheeler A. A review of anatomy, physiology, and benign pathology of the nipple. *Ann Surg Oncol* 2015; 22: 3236–40.
  69. Doucet S, Soussignan R, Sagot P, Schaal B. The secretion of areolar (Montgomery's) glands from lactating women elicits selective, unconditional responses in neonates. *PLoS ONE* 2009; 4: e7579.
  70. Varendi H, Porter RH. Breast odour as the only maternal stimulus elicits crawling towards the odour source. *Acta Paediatr Oslo Nor* 1992; 2001: 372–5.
  71. Varendi H, Christensson K, Porter RH, Winberg J. Soothing effect of amniotic fluid smell in newborn infants. *Early Hum Dev* 1998; 51: 47–55.
  72. Schaal B, Marlier L, Soussignan R. Human fetuses learn odours from their pregnant mother's diet. *Chem Senses* 2000; 25: 729–37.
  73. Mennella JA, Jagnow CP, Beauchamp GK. Prenatal and postnatal flavor learning by human infants. *Pediatrics* 2001; 107: E88.
  74. Delaunay-El Allam M, Soussignan R, Patris B, Marlier L, Schaal B. Long-lasting memory for an odor acquired at the mother's breast: Memory for odor acquired at the mother's breast. *Dev Sci* 2010; 13: 849–63.
  75. Mennella JA. Flavour Programming during Breast-Feeding. In: Goldberg G, Prentice A, Prentice A, Filteau S, Simondon K, editors. *Breast-feeding: early influences on later health*. Dordrecht, the Netherlands: Springer, 2009: 113–20. Available at: [http://link.springer.com/10.1007/978-1-4020-8749-3\\_9](http://link.springer.com/10.1007/978-1-4020-8749-3_9)
  76. Todrank J, Heth G, Restrepo D. Effects of in utero odorant exposure on neuroanatomical development of the olfactory bulb and odour preferences. *Proc R Soc B Biol Sci* 2011; 278: 1949–55.
  77. Einspieler C, Marschik PB, Prechtel HF. Prenatal origin and early postnatal development. *J Psychol* 2008; 216: 148–54.
  78. Prechtel. Prenatal and early postnatal development of human motor behavior. In: Kalverboer AF, Gramsbergen AA, editors. *Handbook of brain and behavior in human development*. Dordrecht, the Netherlands: Kluwer, 2001: 415–28.
  79. Marlier L, Schaal B, Soussignan R. Bottle-fed neonates prefer an odor experienced in utero to an odor experienced postnatally in the feeding context. *Dev Psychobiol* 1998; 33: 133–45.
  80. Tellström R. *Sommar & Vinter i P1*. Sweden: Sveriges Radio, 2017.
  81. DeCasper AJ, Fifer WP. Of human bonding: newborns prefer their mothers' voices. *Science* 1980; 208: 1174–6.
  82. Moon C, Lagercrantz H, Kuhl PK. Language experienced *in utero* affects vowel perception after birth: a two-country study. *Acta Paediatr* 2013; 102: 156–60.
  83. Fawcett C, Arslan M, Falck-Ytter T, Roeyers H, Gredebäck G. Human eyes with dilated pupils induce pupillary contagion in infants. *Sci Rep* 2017; 7. Available from: <http://www.nature.com/articles/s41598-017-08223-3>
  84. Dani C, Cecchi A, Commare A, Rapisardi G, Breschi R, Pratesi S. Behavior of the Newborn during Skin-to-Skin. *J Hum Lact* 2015; 31: 452–7.
  85. Tambini A, Ketz N, Davachi L. Enhanced brain correlations during rest are related to memory for recent experiences. *Neuron* 2010; 65: 280–90.
  86. Cantrill RM, Creedy DK, Cooke M, Dykes F. Effective suckling in relation to naked maternal-infant body contact in the first hour of life: an observation study. *BMC Pregnancy Childbirth* 2014; 14: 20.
  87. Svensson KE, Velandia MI, Matthiesen A-ST, Welles-Nyström BL, Widström A-ME. Effects of mother-infant skin-to-skin contact on severe latch-on problems in older infants: a randomized trial. *Int Breastfeed J* 2013; 8: 1.
  88. Cato K, Sylvén SM, Skalkidou A, Rubertsson C. Experience of the first breastfeeding session in association with the use of the hands-on approach by healthcare professionals: a population-based Swedish study. *Breastfeed Med Off J Acad Breastfeed Med* 2014; 9: 294–300.
  89. Fletcher D, Harris H. The implementation of the HOT program at the Royal Women's Hospital. *Breastfeed Rev Prof Publ Nurs Mothers Assoc Aust* 2000; 8: 19–23.
  90. Kremer KP, Kremer TR. Breastfeeding is associated with decreased childhood maltreatment. *Breastfeed Med* 2018; 13: 18–22.
  91. Christensson K, Siles C, Moreno L, Belaustequi A, De La Fuente P, Lagercrantz H, et al. Temperature, metabolic adaptation and crying in healthy full-term newborns cared for skin-to-skin or in a cot. *Acta Paediatr Oslo Nor* 1992; 1992: 488–93.
  92. Parker LA, Sullivan S, Krueger C, Mueller M. Association of timing of initiation of breastmilk expression on milk volume and timing of lactogenesis stage II among mothers of very low-birth-weight infants. *Breastfeed Med Off J Acad Breastfeed Med* 2015; 10: 84–91.
  93. Pejovic NJ, Herlenius E. Unexpected collapse of healthy newborn infants: risk factors, supervision and hypothermia treatment. *Acta Paediatr* 2013; 102: 680–8.
  94. Poets A, Urschitz MS, Steinfeldt R, Poets CF. Risk factors for early sudden deaths and severe apparent life-threatening events: Table 1. *Arch Dis Child - Fetal Neonatal Ed* 2012; 97: F395–7.
  95. Ludington-Hoe SM, Morgan K. Infant assessment and reduction of sudden unexpected postnatal collapse risk during skin-to-skin contact. *Newborn Infant Nurs Rev* 2014; 14: 28–33.
  96. Becher J-C, Bhushan SS, Lyon AJ. Unexpected collapse in apparently healthy newborns – a prospective national study of a missing cohort of neonatal deaths and near-death events. *Arch Dis Child - Fetal Neonatal Ed* 2012; 97: F30–4.
  97. Matthijse PR, Semmekrot BA, Liem KD. Skin to skin contact and breast-feeding after birth: not always without risk!. *Ned Tijdschr Geneesk* 2016; 160: D171.
  98. Meek JY, Noble L. Implementation of the ten steps to successful breastfeeding saves lives. *JAMA Pediatr* 2016; 170: 925–6.
  99. Bass JL, Gartley T, Kleinman R. Unintended consequences of current breastfeeding initiatives. *JAMA Pediatr* 2016; 170: 923–4.

100. Lehtonen L, Gimeno A, Parra-Llorca A, Vento M. Early neonatal death: A challenge worldwide. *Semin Fetal Neonatal Med* 2017; 22: 153–60.
101. Bartick M, Feldman-Winter L. Skin-to-skin care cannot be blamed for increase in suffocation deaths. *J Pediatr* 2018. Available at: <https://linkinghub.elsevier.com/retrieve/pii/S0022347618307790>
102. Chu DM, Ma J, Prince AL, Antony KM, Seferovic MD, Aagaard KM. Maturation of the infant microbiome community structure and function across multiple body sites and in relation to mode of delivery. *Nat Med* 2017; 23: 314–26.
103. Indrio F, Martini S, Francavilla R, Corvaglia L, Cristofori F, Mastrolia SA, et al. Epigenetic matters: the link between early nutrition, microbiome, and long-term health development. *Front Pediatr* 2017; 5. Available at: <http://journal.frontiersin.org/article/10.3389/fped.2017.00178/full>
104. Schlinzig T, Johansson S, Gunnar A, Ekström T, Norman M. Epigenetic modulation at birth - altered DNA-methylation in white blood cells after Caesarean section. *Acta Paediatr* 2009; 98: 1096–9.
105. Dumas L, Lepage M, Bystrova K, Matthiesen A-S, Welles-Nyström B, Widström A-M. Influence of skin-to-skin contact and rooming-in on early mother-infant interaction: a randomized controlled trial. *Clin Nurs Res* 2013; 22: 310–36.
106. Bystrova K, Ivanova V, Edhborg M, Matthiesen A-S, Ransjö-Arvidson A-B, Mukhamedrakhimov R, et al. Early contact versus separation: effects on mother-infant interaction one year later. *Birth Berkeley Calif* 2009; 36: 97–109.
107. Moffitt TE, Arseneault L, Belsky D, Dickson N, Hancox RJ, Harrington H, et al. A gradient of childhood self-control predicts health, wealth, and public safety. *Proc Natl Acad Sci U S A* 2011; 108: 2693–8.